

basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

MATHEMATICS P2

NOVEMBER 2011

POSSIBLE ANSWERS


MARKS: 150

This memorandum consists of 22 pages.

NOTE:

- If a candidate answers a question TWICE, only mark the FIRST attempt.
- If a candidate has crossed out an attempt of a question and not redone the question, mark the crossed out version.
- Consistent accuracy applies in **ALL** aspects of the marking memorandum.
- Assuming answers/values in order to solve a problem is not acceptable.

QUESTION 1

1.1	Median = 42	✓ answer (1)
1.2	Lower quartile = 32 Upper quartile = 46 Inter quartile range = $46 - 32 = 14$	<p style="border: 1px solid black; padding: 5px;">Answer only: FULL MARKS</p>
1.3	 <p>A box-and-whisker plot on a number line from 20 to 70. The median is at 42. The upper whisker extends to 62, and there is one outlier at 27.</p>	✓ box-and-whisker with a median ✓ skewness ✓ indicating <u>5 number summary</u> 27; 32; 42; 46; 62 or correct scale (3)
1.4	<p>There is a greater spread of scores to the right of the median (42).</p> <p>OR</p> <p>There is a greater spread of scores in the top 50%.</p> <p>OR</p> <p>The spread of the scores on the left hand side of the median is closer to each other.</p> <p>OR</p> <p>The greatest spread of scores lies between Q_3 and the maximum value.</p> <p>Note:</p> <ul style="list-style-type: none"> • Description about the spread based on the box-and-whisker diagram must be accepted. • If it is indicated that it is skewed to the left because the mean is less than the median: full marks 	✓ greater spread ✓ right of median (42) (2) ✓ greater spread ✓ top 50% (2) ✓ spread closer ✓ left of median (2) ✓ greater spread ✓ between Q_3 and max (2) [9]

QUESTION 2

2.1	$\text{Mean} = \frac{\sum_{i=1}^n x_i}{n} = \frac{580}{8} = 72,5$ <p>Note: If rounded off to 73: 1 mark</p>	Answer only: FULL MARKS	✓ 580 ✓ answer (2)
2.2	Standard deviation (σ) = 2,78 (2,783882181...) Note: If rounded off to 2,8: 1 mark		✓✓ answer (2)
2.3	$\therefore 2 \text{ golfers' scores lie outside 1 standard deviation of the mean.}$ <p>The interval for 1 standard deviation of the mean is $(72,5 - 2,78 ; 72,5 + 2,78) = (69,72 ; 75,28)$</p>	Answer only: FULL MARKS	✓ interval ✓ number (2) [6]

QUESTION 3

3.1	30	✓ 30 (1)
3.2	Linear, the points seem to form a straight line.	✓ linear ✓ reason (2)
3.3	<p>The greater the number of hours spent watching TV, the lower the test scores</p> <p>OR</p> <p>The less time a person spends watching TV, the higher the test score.</p> <p>OR</p> <p>Negative correlation between the variables</p> <p>OR</p> <p>Indirect relationship between the variables</p>	✓ deduction (1)
3.4	60 marks. (Accept 50 -70 marks)	✓✓ deduction (2) [6]

QUESTION 4

4.1


TIME	FREQUENCY	CUMULATIVE FREQUENCY
$1 \leq t < 3$	3	3
$3 \leq t < 5$	6	9
$5 \leq t < 7$	7	16
$7 \leq t < 9$	8	24
$9 \leq t < 11$	5	29
$11 \leq t < 13$	1	30

Note: Only cumulative frequency column – full marks

One mark for every two correct cumulative frequency values

(3)

4.2



✓ upper limit
✓ cumulative frequency (at least 4 of 6 y-values correctly plotted)

✓ grounding at (1 ; 0)

✓ shape (not joined by a ruler; smooth curve)

(4)

4.3


Estimated number of learners that took less than 4 minutes:
approximately 5 learners (Accept 6)
Approximate percentage = 16,67% (Accept 20%)

✓ 5 learners
✓ 16,67%

(2)
[9]**Note:**

If using 9 learners and approximate percentage = 30%: 1 mark

If using 5,5 learners and approximate percentage = 18,33%: 1 mark

QUESTION 5

<p>5.1</p> $m_{PQ} \times m_{QR} = -1$ $\left(\frac{a-0}{4+4}\right)\left(\frac{a-0}{4-6}\right) = -1$ $\left(\frac{a}{8}\right)\left(\frac{a}{-2}\right) = -1$ $\frac{a^2}{-16} = -1$ $a^2 = 16$ $a = \pm 4$ $a = 4; \text{ since } a > 0$	<p>OR</p> $PQ^2 + QR^2 = PR^2$ $(8^2 + a^2) + (a^2 + 2^2) = 10^2$ $\therefore 2a^2 = 32$ $\therefore a^2 = 16$ $\therefore a = 4$ <p>OR</p> <p>Let A be the midpoint of diagonal PR.</p> <p>Then $A\left(\frac{-4+6}{2}; \frac{0+0}{2}\right) = A(1; 0)$.</p> <p>$AQ = AR$ (diagonals equal and bisect each other)</p> $AQ^2 = AR^2$ $(1-4)^2 + (0-a)^2 = 5^2$ $9 + a^2 = 25$ $a^2 = 16$ $a = 4$ <p>Note: If candidate uses $a = 4$ at the beginning, then zero marks.</p>	<p>✓ $\frac{a-0}{4+4}$ or $\frac{a}{8}$</p> <p>✓ $\frac{a-0}{4-6}$ or $\frac{a}{-2}$</p> <p>✓ using gradient of perpendicular lines</p> <p>✓ $a^2 = 16$ (4)</p> <p>✓ using Pythagoras</p> <p>✓ $(8^2 + a^2) + (a^2 + 2^2)$</p> <p>✓ 10^2</p> <p>✓ $a^2 = 16$ (4)</p> <p>✓ $(1; 0)$ is centre</p> <p>✓ $AQ = AR$</p> <p>✓ $3^2 + a^2 = 5^2$</p> <p>✓ $a^2 = 16$ (4)</p>
---	--	--

5.2	<p>Equation of line SR:</p> $m_{PQ} = \frac{4 - 0}{4 - (-4)} = \frac{1}{2}$ $m_{SR} = m_{PQ} = \frac{1}{2} \quad PQ \parallel SR$ $y - y_1 = m(x - x_1)$ $y - 0 = \frac{1}{2}(x - 6)$ $y = \frac{1}{2}x - 3$ <p style="text-align: center;">OR</p>	$\checkmark m_{PQ} = \frac{1}{2}$ $\checkmark m_{SR} = \frac{1}{2}$ \checkmark substitution of m and (6 ; 0) \checkmark standard form (4)
-----	---	---

	$m_{PQ} = \frac{1}{2}$ $m_{PQ} = m_{SR} = \frac{1}{2} \quad PQ \parallel SR$ $y = \frac{1}{2}x + c$ $0 = \left(\frac{1}{2}\right)\left(\frac{6}{1}\right) + c$ $-3 = c$ $y = \frac{1}{2}x - 3$ <p style="text-align: center;">OR</p> <p>S(-2 ; -4) (translation)</p> $m_{RS} = \frac{0 + 4}{6 + 2} = \frac{1}{2}$ $\therefore y + 4 = \frac{1}{2}(x + 2)$ $\therefore y = \frac{1}{2}x - 3$	$\checkmark m_{PQ} = \frac{1}{2}$ $\checkmark m_{SR} = \frac{1}{2}$ \checkmark substitution of m and (6 ; 0) \checkmark standard form $\checkmark S(-2 ; -4)$ $\checkmark m_{SR} = \frac{1}{2}$ \checkmark substitution of m and (-2 ; -4) \checkmark standard form (4)
--	---	---

5.3	<p>Eq. of RS: $y = \frac{1}{2}x - 3$</p> <p>Eq. of SP: $y - 0 = -2(x + 4)$</p> $\therefore \frac{1}{2}x - 3 = -2(x + 4)$ $\therefore x = -2$ $y = -4$ <p style="text-align: center;">OR</p>	<p>Answer only: FULL MARKS</p> $\checkmark m = -2$ \checkmark eq. of SP \checkmark value of x \checkmark value of y (4)
-----	--	--

NSC

$$\text{Midpoint PR} = M\left(\frac{-4+6}{2}; \frac{0+0}{2}\right) = (1; 0)$$

Let $S(x; y)$. Then since $M(1; 0)$ is this, the midpoint of QS is:

$$\begin{aligned} \frac{x_1 + x_2}{2} &= 1 & \frac{y_1 + y_2}{2} &= 0 \\ \therefore \frac{x+4}{2} &= 1 & \text{and} & \frac{y+4}{2} = 0 \\ x+4 &= 2 & y+4 &= 0 \\ x &= -2 & y &= -4 \end{aligned}$$

OR

$$\sqrt{\frac{x+4}{2}} = 1$$

$$\sqrt{\frac{y+4}{2}} = 0$$

✓ value of x ✓ value of y

(4)

The translation that sends $Q(4; 4)$ to $R(6; 0)$ also sends $P(-4; 0)$ to S .

$$(6; 0) = (4+2; 4-4)$$

$$\therefore S = (-4+2; 0-4) = (-2; -4)$$

- ✓ method
- ✓ 2 or $x+2$
- ✓ -4 or $y-4$
- ✓ answer

(4)

OR

The translation that sends $Q(4; 4)$ to $P(-4; 0)$ also sends $R(6; 0)$ to S .

$$(-4; 0) = (4-8; 4-4)$$

$$\therefore S = (6-8; 0-4) = (-2; -4)$$

- ✓ method
- ✓ -8 or $x-8$
- ✓ -4 or $y-4$
- ✓ answer

(4)

OR

$$m_{PQ} = m_{SR}$$

$$\frac{1}{2} = \frac{y}{x-6}$$

$$2y = x-6 \quad (1)$$

✓ equations using the gradient

$$m_{PS} = m_{SR}$$

$$\frac{y}{x+4} = \frac{4}{-2}$$

$$-2y = 4x + 16 \quad (2)$$

✓ adding the equations

$$(1) + (2); 0 = 5x + 10$$

$$x = -2$$

$$\text{Substitute: } 2y = -2 - 6 = -8$$

$$y = -4$$

- ✓ value of x
- ✓ value of y

(4)

5.4

$$PR = 6 - (-4)$$

$$= 10$$

OR

$$PR^2 = (6+4)^2 + (0-0)^2$$

$$PR = 10$$

Answer only:
FULL MARKS


$$\checkmark 6 - (-4)$$

$$\checkmark 10$$

(2)

- ✓ substitution in correct formula
- ✓ 10

		NSC -	
5.5	$\text{midpoint PR} = \left(\frac{6+(-4)}{2}, \frac{0+0}{2} \right) = (1; 0)$ $\text{radius of circle} = \frac{1}{2} \text{PR} = 5 \text{ units}$ $\therefore (x-1)^2 + (y-0)^2 = 5^2$ $(x-1)^2 + y^2 = 25$	Answer only: FULL MARKS	✓ midpoint ✓ radius ✓ eq. of circle in correct form (3)
5.6	$(x-1)^2 + y^2 = 25$ substitute Q(4 ; 4): $\text{LHS} = (4-1)^2 + 4^2$ $= 25$ $= \text{RHS}$ $\therefore Q$ is a point on the circle Note: If substitute point into equation resulting in $25 = 25$: 1 mark No conclusion: 1 mark	OR	✓ substitute Q(4;4) ✓ LHS = RHS (2)
	Distance from centre (1 ; 0) to Q(4 ; 4) $\therefore Q$ is a point on circle, $r = 5$ OR PR is the diameter of circle PQR therefore Q lies on circle ($P\hat{Q}R = 90^\circ$)	OR	✓ = 5 ✓ conclusion (2) ✓ diameter ✓ $P\hat{Q}R = 90^\circ$ (2)
	$(4-1)^2 + y^2 = 25$ $y^2 = 16$ $\therefore y = 4$ $\therefore Q$ is a point on the circle	OR	✓ substitute $x = 4$ ✓ conclusion (2)
	$(x-1)^2 + 4^2 = 25$ $(x-1)^2 = 9$ $x-1 = 3$ $x = 4$ $\therefore Q$ is a point on the circle	OR	✓ substitute $y = 4$ ✓ conclusion (2)
5.7	P needs to shift at least 4 units to the right and S needs to shift at least 4 units up for the image of PQRS in first quadrant. \therefore minimum value of k is 4 and minimum value of l is 4 \therefore minimum value of $k + l$ is 8	Answer only: FULL MARKS	✓ $k = 4$ ✓ $l = 4$ ✓ $k + l = 8$ (3) [22]
	Note: No CA mark applies in 5.7 if k and l are not minimums.		

QUESTION 6

6.1	$x_C = x_B = -1$ $y_C = y_B + 5 = 6$ $\therefore C(-1; 6)$	✓ value of x ✓ value of y (2)
6.2	$BA \perp CA$ (tangent \perp radius) $\therefore CA^2 = BC^2 - AB^2$ (Pythagoras) $= (5)^2 - (\sqrt{20})^2 = 5$ $\therefore CA = \sqrt{5}$ or 2,24 units	✓ $BA \perp CA$ or $\hat{BAC} = 90^\circ$ ✓ substitution into Pythagoras ✓ answer (3)
6.3	$\tan \theta = \frac{\sqrt{5}}{\sqrt{20}} = \frac{\sqrt{5}}{2\sqrt{5}} = \frac{1}{2}$	✓ tan ratio (in any form) (1)
6.4	$m_{DC} \times m_{AB} = -1$ $m_{DC} = \tan \theta = \frac{1}{2}$ $m_{DC} = \frac{1}{2}$ $m_{AB} = -2$	✓ $m_{DC} \times m_{AB} = -1$ ✓ $m_{DC} = \tan \theta = \frac{1}{2}$ (2)

NSC -

<p>6.5</p> <p>Eq. of DC: $y - 6 = \frac{1}{2}(x + 1)$</p> $y = \frac{1}{2}x + \frac{13}{2}$ <p>Eq. of AB: $y - 1 = -2(x + 1)$</p> $y = -2x - 1$ $-2x - 1 = \frac{1}{2}x + \frac{13}{2}$ $-\frac{5}{2}x = \frac{15}{2}$ $x = -3$ $y = -2(-3) - 1$ $y = 5$ $\therefore A(-3 ; 5)$	<p>Answer only: $(-3 ; 5)$: 1 mark</p> <p>✓ DC: subst m and $(-1 ; 6)$ ✓ eq. of DC</p> <p>✓ eq. of AB</p> <p>✓ equating equations</p> <p>✓ value of x ✓ value of y</p> <p style="text-align: right;">(6)</p>
<p style="text-align: center;">OR</p> <p>Eq. of DC: $y - 6 = \frac{1}{2}(x + 1)$</p> $y = \frac{1}{2}x + \frac{13}{2}$ <p>Eq. of AB: $y - 1 = -2(x + 1)$</p> $y = -2x - 1$ <p><u>At A:</u> $x - 2(-2x - 1) + 13 = 0$ $x + 4x + 2 + 13 = 0$ $5x = -15$ $x = -3$ and $y = -2(-3) - 1 = 5$ $\therefore A(-3 ; 5)$</p>	<p>✓ DC: subst m and $(-1 ; 6)$ ✓ eq. of DC</p> <p>✓ subt m and $(-1 ; 1)$ ✓ eq. of AB</p> <p>✓ value of x ✓ value of y</p> <p style="text-align: right;">(6)</p>

<p>Eq. of DC: $y - 6 = \frac{1}{2}(x + 1)$</p> $y = \frac{1}{2}x + \frac{13}{2}$ <p>Eq. of circle: $(x + 1)^2 + (y - 1)^2 = 20$</p> <p><u>At A:</u></p> $(x + 1)^2 + (\frac{1}{2}x + \frac{13}{2} - 1)^2 = 20$ $(x + 1)^2 + (\frac{1}{2}x + \frac{11}{2})^2 = 20$ $1\frac{1}{4}x^2 + \frac{15}{2}x + 11\frac{1}{4} = 0$ $\therefore x^2 + 6x + 9 = 0$ $(x + 3)^2 = 0$ $\therefore x = -3$ <p>and $y = \frac{1}{2}(-3) + \frac{13}{2} = 5$</p> $\therefore A(-3 ; 5)$	<p>✓ DC: subst m and $(-1 ; 6)$ ✓ eq. of DC</p> <p>✓ substitution</p> <p>$x^2 + 6x + 9 = 0$</p> <p>✓ value of x</p> <p>✓ value of y</p> <p style="text-align: right;">(6)</p>
---	--

NSC -

ORDraw $AE \perp BC$

$$\cos \theta = \frac{2\sqrt{5}}{5} = \frac{AE}{\sqrt{5}} = \frac{BE}{2\sqrt{5}}$$


$$\therefore AE = \frac{2 \times 5}{5} = 2$$

$$BE = \frac{4 \times 5}{5} = 4$$

$$x_A = -1 - AE = -1 - 2 = -3$$

$$\therefore y_A = 1 + BE = 4 + 1 = 5$$

$$\therefore A(-3; 5)$$



$$\checkmark \frac{2\sqrt{5}}{5} = \frac{AE}{\sqrt{5}}$$

$$\checkmark AE = 2$$

$$\checkmark \frac{2\sqrt{5}}{5} = \frac{BE}{2\sqrt{5}}$$

$$\checkmark BE = 1$$

$$\checkmark -3$$

$$\checkmark 5$$

(6)

OR

$$(x+1)^2 + (y-1)^2 = 20 \quad (1)$$

$$y = -2x - 1 \quad (2)$$

$$(x+1)^2 + (-2x-2)^2 = 20$$

$$x^2 + 2x + 1 + 4x^2 + 8x + 4 - 20 = 0$$

$$5x^2 + 10x - 15 = 0$$

$$x^2 + 10x - 15 = 0$$

$$(x+3)(x-1) = 0$$

$$x = -3 \text{ or } x \neq 1$$

subst (1) in (2)

$$\therefore y = 5$$

 \checkmark subst m and
 $(-1; 1)$
 \checkmark eq of AB

 \checkmark eq of circle


 \checkmark substation

 \checkmark value of x
 \checkmark value of y (6)

NSC - 1

OR

$$\text{Equation AC : } y = \frac{1}{2}x + 6 \frac{1}{2}$$



$$\tan \theta = \frac{1}{2}$$

$$\theta = 26.57^\circ$$

$$AP = \sqrt{5} \cos 26.57^\circ$$

$$AP = 2$$

$$CP = \sqrt{5} \sin 26.57^\circ$$

$$CP = 1$$

$$\therefore x = -1 - 2 = -3$$

$$y = 6 - 1 = 5$$

$$\therefore A(-3; 5)$$

$$\checkmark \theta = 26.57^\circ$$

✓

$$AP = \sqrt{5} \cos 26.57^\circ$$

$$\checkmark AP = 2$$

$$\checkmark CP = 1$$

$$\checkmark \text{ value of } x$$

$$\checkmark \text{ value of } y$$

(6)

6.6

$$\text{Area } \Delta ABC = \frac{1}{2}(\sqrt{5})(\sqrt{20}) = 5$$

$$\checkmark \frac{1}{2}(\sqrt{5})(\sqrt{20})$$

$$\text{Eqn. of DC is } y = \frac{1}{2}x + \frac{13}{2}$$

$$\checkmark OF = \frac{13}{2}$$

$$\text{Therefore } OF = \frac{13}{2} \text{ and } OD = 13.$$

$$\checkmark OD = 13$$

$$\text{Area } \Delta ODF = \frac{1}{2}\left(\frac{13}{2}\right)(13) = \frac{169}{4}$$

$$\checkmark \frac{1}{2}\left(\frac{13}{2}\right)(13)$$

$$\text{Area } \Delta ABC : \text{Area } \Delta ODF = 5 : \frac{169}{4} = 20 : 169$$

✓ answer

(5)

OR

$$DF^2 = 13^2 + \left(\frac{13}{2}\right)^2 = \frac{845}{4}$$

$$\checkmark = 13^2$$

$$DF = \frac{13\sqrt{5}}{2}$$

$$+ \left(\frac{13}{2}\right)^2 = \frac{845}{4}$$

$$\frac{\Delta ABC}{\Delta ODF} = \frac{\frac{1}{2}(5)(\sqrt{20}) \sin \theta}{\frac{1}{2}(13)\left(\frac{13\sqrt{5}}{2}\right) \sin \theta}$$

$$\checkmark DF = \frac{13\sqrt{5}}{2}$$

$$= \frac{20}{169}$$

$$\checkmark \frac{1}{2}(5)(\sqrt{20}) \sin \theta$$

$$= \frac{1}{2}(13)\left(\frac{13\sqrt{5}}{2}\right) \sin \theta$$

✓ answer (5)



OR

ΔODF is an enlargement of ΔABC
 $\therefore \text{area } \Delta ABC : \text{area } \Delta ODF = AB^2 : OD^2 = 20 : OD^2$
 Equation of DC is $y = \frac{1}{2}x + \frac{13}{2}$
 $x_D = -13$
 $OD = 13$
 $\therefore \text{area } \Delta ABC : \text{area } \Delta ODF = AB^2 : OD^2 = 20 : 169$

✓ enlargement
 ✓✓
 $AB^2 : OD^2 = 20 : OD^2$
 ✓ - 13
 ✓ answer (5)

[19]

QUESTION 7

7.1	$(x; y) \rightarrow (x+4; y) \rightarrow (-x-4; -y)$ OR $(x; y) \rightarrow (-x-4; -y)$	✓ $x+4$ ✓ y ✓ $-x-4$ ✓ $-y$ (4)
7.2	New centre = $(-2; -5)$ $(x+2)^2 + (y+5)^2 = 16$ $x^2 + 4x + 4 + y^2 + 10y + 25 - 16 = 0$ $x^2 + y^2 + 4x + 10y + 13 = 0$	✓ $(-2; -5)$ ✓ $(x+2)^2 + (y+5)^2$ ✓ 16 ✓ simplification (4) [8]



QUESTION 8

8.1	Rotation of 90° anticlockwise about the origin. OR Rotation of 270° clockwise about the origin. Note: if reflection of 90° anticlockwise: 0 marks	✓ rotation 90° ✓ anticlockwise (2) ✓ rotation 270° ✓ clockwise (2)
8.2	$D(5; -4)$ $D'(4; 5)$	✓ 4 ✓ 5 (2)
8.3	$G(-7; -6)$	✓ -7 ✓ -6 (2)
8.4	Area ABCD = $5 \times 2 = 10$ square units $\text{Area MNRP} = 10 \times \left(\frac{3}{2}\right)^2 = \frac{45}{2}$ $\text{Area ABCD} \times \text{Area MNRP}$ $= 10 \times \frac{9}{4} \times 10$ $= 225 (\text{units})^4$	✓ area ABCD = 10 ✓ area MNRP $= \frac{45}{2}$ ✓ 225 (3)

OR

	$\text{Product} = \left(\frac{3}{2}\right)^2 \times (\text{area } ABCD)^2$ $= \frac{9}{4} \times (5 \times 2)^2$ $= 225 \text{ (units)}^4$ <p>Note: CA will apply if $\left(\frac{3}{2}\right)^2$ used in calculation.</p>	$\checkmark \left(\frac{3}{2}\right)^2$ $\checkmark 10^2$ $\checkmark 225$ (3) [9]
--	---	--

QUESTION 9

9.1	9.1.1	 or 	\checkmark sketch $\checkmark r = 7$ $\checkmark \frac{\sqrt{40}}{7}$ (3)
	9.1.2	$\sin(180^\circ + A)$ $= -\sin A$ $= -\frac{3}{7}$ <p style="text-align: center;">OR</p> $\begin{aligned} \sin(180^\circ + A) &= \sin 180^\circ \cdot \cos A + \cos 180^\circ \cdot \sin A \\ &= 0 \cdot \cos A - 1 \cdot \sin A \\ &= -\sin A \\ &= -\frac{3}{7} \end{aligned}$	$\checkmark -\sin A$ $\checkmark -\frac{3}{7}$ (2)
9.2		$\begin{aligned} &\frac{\cos 100^\circ \times \tan^2 120^\circ}{\sin(-10^\circ)} \\ &= \frac{(-\cos 80^\circ)(-\tan 60^\circ)^2}{(-\sin 10^\circ)} \\ &= \frac{(-\cos 80^\circ) \times ((-\sqrt{3})^2)}{(-\cos 80^\circ)} \\ &= 3 \end{aligned}$ <p>Note: Answer only: 0 marks</p> <p>Note: If $\frac{+\cos 80^\circ}{+\sin 10^\circ}$ (assume two negatives cancelled), no penalty</p>	$\checkmark -\cos 80^\circ$ $\checkmark -\tan 60^\circ$ or $\tan^2 60^\circ$ $\checkmark -\sin 10^\circ$ $\checkmark -\sqrt{3}$ $\checkmark \sin 10^\circ =$ $\cos 80^\circ$ $\checkmark 3$ (6)

1

NSC -

OR


$$\begin{aligned} & \frac{\cos 100^\circ \times \tan^2 120^\circ}{\sin (-10^\circ)} \\ &= \frac{(-\cos 80^\circ)(-\tan 60^\circ)^2}{(-\sin 10^\circ)} \\ &= \frac{(-\sin 10^\circ) \times ((-\sqrt{3})^2)}{(-\sin 10^\circ)} \\ &= 3 \end{aligned}$$

- ✓ $-\cos 80^\circ$
 - ✓ $-\sin 10^\circ$
 - ✓ $-\tan 60^\circ$
 - ✓ $-\sqrt{3}$
 - ✓ $\cos 80^\circ = \sin 10^\circ$
 - ✓ 3
- (6)

OR

$$\begin{aligned} & \frac{\cos 100^\circ \times \tan^2 120^\circ}{\sin (-10^\circ)} \\ &= \frac{\cos(90^\circ + 10^\circ)}{-\sin(10^\circ)} \times \tan^2 60^\circ \\ &= \frac{-\sin 10^\circ}{-\sin 10^\circ} \times (\sqrt{3})^2 \\ &= 3 \end{aligned}$$

- ✓ $\cos(90^\circ + 10^\circ)$
 - ✓ $-\sin 10^\circ$
 - ✓ $-\sin 10^\circ$
 - ✓ $\tan^2 60^\circ$
 - ✓ $\sqrt{3}$
 - ✓ 3
- (6)



9.3	9.3.1	$r = 5$ $\sin ROP = \frac{3}{5} = 0,6$	✓ 5 ✓ ratio
	9.3.2	$ROP = 36,87^\circ$ $QOP = 180^\circ - 36,869\dots^\circ$ $QOP = 143,13^\circ$	✓ $36,869\dots^\circ$ ✓ $143,13^\circ$

Answer only: Full Marks

	<p>9.3.3</p> $x_m = x \cos \theta + y \sin \theta$ $a = 4 \cos 115^\circ + 3 \sin 115^\circ$ $a = 1,03$	<p>Note: Penalise 1 mark for rounding incorrectly Note: If incorrect angle is used in the x- formula: 1 mark</p>	<ul style="list-style-type: none"> ✓ formula ✓ substitution of values ✓ $a = 1,03$ (3)
	<p>OR</p> <p>Rotation of 115° clockwise = 245° anticlockwise</p> $x_m = x \cos \theta - y \sin \theta$ $a = 4 \cos 245^\circ - 3 \sin 245^\circ$ $a = 1,03$	<ul style="list-style-type: none"> ✓ formula ✓ substitution of values ✓ $a = 1,03$ (3)	

OR

$$\tan P\hat{O}R = \frac{3}{4}$$

$$P\hat{O}R = 36,86\dots^\circ$$

$$M\hat{O}R = 78,13\dots^\circ$$

$$\cos M\hat{O}R = \frac{a}{5}$$

$$a = 5 \cos 78,13^\circ$$

$$a = 1,03$$

$$\checkmark 36,86^\circ$$

$$\checkmark \cos \text{ratio}$$

$$\checkmark a = 1,03$$

(3)

[18]**QUESTION 10**

10.1	$f(225^\circ) = 2$ $\therefore a \tan 225^\circ = 2 \quad \therefore a = 2$ $g(0) = 4$ $\therefore b \cos 0^\circ = 4 \quad \therefore b = 4$	<p>Answer only: Full marks</p>	<ul style="list-style-type: none"> ✓ substitution ✓ $a = 2$ ✓ substitution ✓ $b = 4$ (4)
10.2	Minimum value of $g(x) + 2 = -4 + 2 = -2$	<p>Answer only: Full marks</p>	<ul style="list-style-type: none"> ✓ -4 ✓ -2 (2)
10.3	$\text{Period} = \frac{180^\circ}{\frac{1}{2}} = 360^\circ$	<p>Answer only: Full marks</p>	<ul style="list-style-type: none"> ✓ $\frac{180^\circ}{\frac{1}{2}}$ ✓ 360° (2)

<p>10.4 At P $f(\theta) = g(\theta)$</p> $2\tan \theta = 4\cos \theta$ <p>for $180^\circ - \theta$: $2\tan(180^\circ - \theta) = -2\tan \theta$ and $4\cos(180^\circ - \theta) = -4\cos \theta$</p> <p>$2\tan \theta = 4\cos \theta$ at P</p> $\therefore -2\tan \theta = -4\cos \theta$ $\therefore 2\tan(180^\circ - \theta) = 4\cos(180^\circ - \theta)$ at Q	$\checkmark 2\tan \theta = 4\cos \theta$ $\checkmark 2\tan(180^\circ - \theta) = -2\tan \theta$ $\checkmark 4\cos(180^\circ - \theta) = -4\cos \theta$ $\checkmark 2\tan(180^\circ - \theta) = 4\cos(180^\circ - \theta)$ (4)
<p>OR</p> $2\tan \theta = 4\cos \theta$ $\frac{\sin \theta}{\cos \theta} = 2\cos \theta$ $\sin \theta = 2\cos^2 \theta$ $= 2(1 - \sin^2 \theta)$ $2\sin^2 \theta + \sin \theta - 2 = 0$ $\sin \theta = \frac{-1 \pm \sqrt{1 - 4(2)(-2)}}{4}$ $\sin \theta = 0,78077\dots$ $\theta = 51,33^\circ \text{ or } 128,67^\circ$ $\therefore \text{the } x\text{-coordinate of Q is } 180^\circ - x_p$	\checkmark equation $\checkmark \sin \theta = 0,78077\dots$ $\checkmark 51,33^\circ$ $\checkmark 128,67^\circ$ (4) [12]

QUESTION 11

<p>11.1 Area $\Delta ABC = \frac{1}{2} \cdot AB \cdot BC \cdot \sin 50^\circ$</p> $= \frac{1}{2}(5)(5)\sin 50^\circ$ $= 9,58 \text{ units}^2$	\checkmark substitution into correct formula \checkmark answer (2)
<p>OR</p> <p>Area of ΔABC</p> $= \frac{1}{2}(2)(5)\sin 25^\circ(5\cos 25^\circ)$ $= 9,58 \text{ units}^2$	\checkmark base and height in terms of 5 and 25° \checkmark answer (2)
<p>OR</p> <p>Area of ΔABC</p> $= [\frac{1}{2}(5\cos 65^\circ)(5\sin 65^\circ)](2)$ $= 9,58 \text{ units}^2$	\checkmark base and height in terms of 5 and 65° \checkmark answer (2)

11.2	$AC^2 = 5^2 + 5^2 - 2(5)(5)\cos 50^\circ$ $AC^2 = 17,86061952$ $AC = 4,23 \text{ units}$ <p style="text-align: center;">OR</p>	<ul style="list-style-type: none"> ✓ use of cosine rule ✓ substitution ✓ answer (3)
	$\hat{A} = \hat{C} = 65^\circ \quad (\text{angles opposite equal sides})$ $\frac{\sin 65^\circ}{5} = \frac{\sin 50^\circ}{AC}$ $AC = \frac{5 \sin 50^\circ}{\sin 65^\circ}$ $= 4,23 \text{ units}$ <p style="text-align: center;">OR</p>	<ul style="list-style-type: none"> ✓ use of sine rule ✓ substitution ✓ answer (3)
	$\sin 25^\circ = \frac{\frac{1}{2}(AC)}{5}$ $AC = 2(5) \sin 25^\circ$ $= 4,23 \text{ units}$ <p style="text-align: center;">OR</p> $\cos 65^\circ = \frac{\frac{1}{2}(AC)}{5}$ $AC = 2(5) \cos 65^\circ$ $AC = 4,23 \text{ units}$	<ul style="list-style-type: none"> ✓ sketch/diagram ✓ $\sin 25^\circ = \frac{1}{2} AC$ ✓ answer (3)
11.3	$\tan 25^\circ = \frac{CF}{AC}$ $\therefore CF = 4,23 \times \tan 25^\circ$ $\therefore CF = 1,97 \text{ units}$ <p style="text-align: center;">OR</p> $\frac{FC}{\sin 25^\circ} = \frac{4,23}{\sin 65^\circ}$ $FC = \frac{4,23 \sin 25^\circ}{\sin 65^\circ}$ $= 1,97 \text{ units}$	<ul style="list-style-type: none"> ✓ ratio ✓ CF as subject ✓ answer (3) <ul style="list-style-type: none"> ✓ sine rule ✓ FC as subject ✓ answer (3)

[8]

QUESTION 12

12.1	$ \begin{aligned} LHS &= \frac{\sin(360^\circ + 90^\circ + x - \alpha)}{\cos(\alpha - x)} \\ &= \frac{\sin(90^\circ + x - \alpha)}{\cos(\alpha - x)} \\ &= \frac{\cos(x - \alpha)}{\cos(\alpha - x)} \\ &= \frac{\cos(\alpha - x)}{\cos(\alpha - x)} \\ &= 1 \end{aligned} $	<ul style="list-style-type: none"> ✓ subtracting 360° ✓ cos (x - α) ✓ cos(α - x) <p>(3)</p>
	OR	
	$ \begin{aligned} LHS &= \frac{\sin[90^\circ - (\alpha - x)]}{\cos(\alpha - x)} \\ &= \frac{\cos(\alpha - x)}{\cos(\alpha - x)} \\ &= 1 \\ &= RHS \end{aligned} $	<ul style="list-style-type: none"> ✓ subtracting 360° ✓ writing as 90° - (α - x) ✓ cos(α - x) <p>(3)</p>
12.2	$ \begin{aligned} \cos 2x &= 1 - 3 \cos x \\ 2 \cos^2 x - 1 &= 1 - 3 \cos x \\ 2 \cos^2 x + 3 \cos x - 2 &= 0 \\ (2 \cos x - 1)(\cos x + 2) &= 0 \\ \cos x = \frac{1}{2} &\quad \text{or } \cos x = -2 \\ &\quad \text{n/a} \\ x = 60^\circ + k \cdot 360^\circ ; k \in \mathbb{Z} &\quad \text{or } x = 300^\circ + k \cdot 360^\circ ; k \in \mathbb{Z} \end{aligned} $	<ul style="list-style-type: none"> ✓ $\cos 2x = 2 \cos^2 x - 1$ ✓ factorisation ✓ $\cos x = \frac{1}{2}$ ✓ 60° ✓ 300° ✓ + k.360° ✓ k ∈ Z <p>(7)</p>
	OR	
	$x = \pm 60^\circ + k \cdot 360^\circ ; k \in \mathbb{Z}$	
12.3.1	$ \begin{aligned} \text{LHS:} \\ &\frac{\sin A \cos B - \cos A \sin B}{\sin B \cos B} \\ &= \frac{\sin(A - B)}{\sin B \cos B} \\ \text{RHS} &= \frac{2 \sin(A - B)}{2 \sin B \cos B} \\ &= \frac{\sin(A - B)}{\sin B \cos B} \\ &= \text{LHS} \end{aligned} $	<ul style="list-style-type: none"> ✓ writing as single fraction ✓ comp. angle expansion ✓ comp. angle expansion ✓ simplification <p>(4)</p>

NSC –
OR

LHS:

$$\begin{aligned}
 & \frac{\sin A \cos B - \cos A \sin B}{\sin B \cos B} \\
 &= \frac{\sin(A - B)}{\sin B \cos B} \\
 &= \frac{2 \sin(A - B)}{2 \sin B \cos B} \\
 &= \frac{2 \sin(A - B)}{\sin 2B} \\
 &= RHS
 \end{aligned}$$

- ✓ writing as single fraction
- ✓ comp. angle expansion
- ✓ mult. by 2
- ✓ comp. angle expansion

(4)

OR

$$\begin{aligned}
 RHS &= \frac{2 \sin(A - B)}{\sin 2B} \\
 &= \frac{2(\sin A \cos B - \cos A \sin B)}{2 \sin B \cos B} \\
 &= \frac{\sin A \cos B - \cos A \sin B}{\sin B \cos B} \\
 &= \frac{\sin A \cos B}{\sin B \cos B} - \frac{\cos A \sin B}{\sin B \cos B} \\
 &= \frac{\sin A}{\sin B} - \frac{\cos A}{\cos B} \\
 &= LHS
 \end{aligned}$$

- ✓ expansion
- ✓ expansion
- ✓ divide by 2
- ✓ write as separate fractions

(4)

12.3.2(a)	$\begin{aligned} A &= 5B \\ \frac{\sin 5B}{\sin B} - \frac{\cos 5B}{\cos B} &= \frac{2 \sin(5B - B)}{\sin 2B} \\ &= \frac{2 \sin 4B}{\sin 2B} \\ &= \frac{4 \sin 2B \cos 2B}{\sin 2B} \\ &= 4 \cos 2B \end{aligned}$ <p style="text-align: center;">OR</p> $\begin{aligned} \frac{\sin 5B}{\sin B} - \frac{\cos 5B}{\cos B} &= \frac{\sin 5B \cos B - \cos 5B \sin B}{\sin B \cos B} \\ &= \frac{\sin(5B - B)}{\sin B \cos B} \\ &= \frac{\sin 4B}{\frac{1}{2}(2) \sin B \cos B} \\ &= \frac{2 \sin 2B \cos 2B}{\frac{1}{2} \sin 2B} \\ &= 4 \cos 2B \end{aligned}$	<ul style="list-style-type: none"> ✓ recognising $A = 5B$ ✓ substituting $A = 5B$ ✓ $\sin 4B$ = $2 \sin 2B \cos 2B$ <p style="text-align: right;">(3)</p> <ul style="list-style-type: none"> ✓ writing as single fraction ✓ $\sin 4B$ = $2 \sin 2B \cos 2B$ ✓ compound angle in denominator <p style="text-align: right;">(3)</p>
12.3.2(b)	$\begin{aligned} B &= 18^\circ \\ \frac{\sin 90^\circ}{\sin 18^\circ} - \frac{\cos 90^\circ}{\cos 18^\circ} &= 4 \cos 2(18)^\circ \\ \therefore \frac{1}{\sin 18^\circ} - 0 &= 4 \cos 36^\circ \\ \therefore \frac{1}{\sin 18^\circ} &= 4 \cos 36^\circ \end{aligned}$	<ul style="list-style-type: none"> ✓ recognising $B = 18^\circ$ ✓ substituting $B = 18^\circ$ ✓ simplify <p style="text-align: right;">(3)</p>
12.3.2(c)	<p>Let $\sin 18^\circ = a$</p> $\begin{aligned} \frac{1}{\sin 18^\circ} &= 4 \cos 36^\circ \\ \frac{1}{\sin 18^\circ} &= 4(1 - 2 \sin^2 18^\circ) \\ \therefore \frac{1}{a} &= 4(1 - 2a^2) \\ \therefore 1 &= 4a - 8a^3 \\ \therefore 8a^3 - 4a + 1 &= 0 \end{aligned}$ <p>Hence $\sin 18^\circ$ is a solution of $\therefore 8x^3 - 4x + 1 = 0$</p> <p style="text-align: center;">OR</p> <p style="text-align: center;">—</p>	<ul style="list-style-type: none"> ✓ $\sin 18^\circ = a$ ✓ $\cos 36^\circ = 1 - 2 \sin^2 18^\circ$ ✓ substitution of a ✓ simplification <p style="text-align: right;">(4)</p>

	$\frac{1}{\sin 18^\circ} = 4 \cos 36^\circ$ $\frac{1}{\sin 18^\circ} = 4(1 - 2 \sin^2 18^\circ)$ $\frac{1}{\sin 18^\circ} = 4 - 8 \sin^2 18^\circ$ $8(\sin 18^\circ)^3 - 4(\sin 18) + 1 = 0$ <p>Hence $\sin 18^\circ$ is a solution of $\therefore 8x^3 - 4x + 1 = 0$</p>	<ul style="list-style-type: none"> ✓ $\cos 36^\circ$ = $1 - 2 \sin^2 18^\circ$ ✓ simplification ✓ equation i.t.o $\sin 18^\circ$ ✓ replacing $\sin 18^\circ = x$ <p>(4) [24]</p>
	Note: substituting $x = \sin 18^\circ$ into $8x^3 - 4x + 1$ using a calculator showing equal to 0: 0 marks	

TOTAL: 150