



# basic education

---

Department:  
Basic Education  
**REPUBLIC OF SOUTH AFRICA**

**SENIOR CERTIFICATE EXAMNATIONS**  
***SENIORSERTIFIKAAT-EKSAMEN***

**MATHEMATICS P2/*WISKUNDE V2***

**2017**

**MARKING GUIDELINES/*NASIENRIGLYNE***

**MARKS: 150**  
***PUNTE: 150***

**These marking guidelines consist of 22 pages.**  
***Hierdie nasienriglyne bestaan uit 22 bladsye..***

**NOTE:**

- If a candidate answers a question TWICE, only mark the FIRST attempt.
- If a candidate has crossed out an attempt of a question and not redone the question, mark the crossed out version.
- Consistent accuracy applies in ALL aspects of the marking memorandum. Stop marking at the second calculation error.
- Assuming answers/values in order to solve a problem is NOT acceptable.
- Geometry:  
S = a mark for a correct statement (a statement mark is independent of a reason)  
R = a mark for a correct reason (a reason mark may only be awarded if the statement is correct)  
S/R = award a mark if statement and reason are both correct

**NOTA:**


- *As 'n kandidaat 'n vraag TWEEKEER beantwoord, merk slegs die EERSTE poging.*
- *As 'n kandidaat 'n antwoord van 'n vraag doodtrek en nie oordoen nie, merk die doodgetrekte poging.*
- *Volgehoue akkuraatheid word in ALLE aspekte van die memorandum toegepas. Hou op nasien by die tweede berekeningsfout.*
- *Aanvaar van antwoorde/waardes om 'n probleem op te los, word NIE toegelaat nie.*
- *Euklidiese Meetkunde:*  
S = 'n punt vir 'n korrekte bewering ('n beweringspunt is onafhanklik van die rede)  
R = 'n punt vir 'n korrekte rede ('n punt kan slegs vir 'n rede toegeken word, indien die bewering korrek is  
S/R = 'n punt word toegeken indien beide die bewering en rede korrek is

**QUESTION/VRAAG/VRAAG 1**


<b>TIME TAKEN (IN HOURS)</b>	5	7	5	8	10	13	15	20	18	25	23
<b>COST (IN THOUSANDS OF RANDS)</b>	10	10	15	12	20	25	28	32	28	40	30

1.1	$a = 4,806\dots = 4,81$ $b = 1,323\dots = 1,32$ $y = 4,81 + 1,32x$	✓ $a = 4,81$ ✓ $b = 1,32$ ✓ equation (3)
1.2	Cost = 25,974... = 25,97 thousand rand (calculator) = R25 970  <b>OR/OF</b>  $y = 4,81 + 1,32(16)$ $y = 25,93$ Cost = R25 930	✓ 25,97 ✓ answer (in Rands) (2)  ✓ substitution  ✓ answer (in Rands) (2)
1.3	$r = 0,949\dots = 0,95$	✓ answer (1)
1.4	$x = 0$ $y = 4,81$ <b>OR</b> (4,80647) $\therefore$ R4 810 <b>OR</b> R4806,47	✓ $x = 0$  ✓ answer (2) <b>[8]</b>

**QUESTION/VRAAG 2**

2.1	modal class: $80 < x \leq 100$	✓ correct class (1)																		
2.2	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 33%;">Commission earned (in thousands of Rands)</th> <th style="width: 33%;">Frequency</th> <th style="width: 33%;">Cumulative Frequency</th> </tr> </thead> <tbody> <tr> <td><math>20 &lt; x \leq 40</math></td> <td>7</td> <td>7</td> </tr> <tr> <td><math>40 &lt; x \leq 60</math></td> <td>6</td> <td>13</td> </tr> <tr> <td><math>60 &lt; x \leq 80</math></td> <td>8</td> <td>21</td> </tr> <tr> <td><math>80 &lt; x \leq 100</math></td> <td>10</td> <td>31</td> </tr> <tr> <td><math>100 &lt; x \leq 120</math></td> <td>4</td> <td>35</td> </tr> </tbody> </table>	Commission earned (in thousands of Rands)	Frequency	Cumulative Frequency	$20 < x \leq 40$	7	7	$40 < x \leq 60$	6	13	$60 < x \leq 80$	8	21	$80 < x \leq 100$	10	31	$100 < x \leq 120$	4	35	<p>✓ 13 ; 21</p> <p>✓ 31 ; 35 (2)</p>
Commission earned (in thousands of Rands)	Frequency	Cumulative Frequency																		
$20 < x \leq 40$	7	7																		
$40 < x \leq 60$	6	13																		
$60 < x \leq 80$	8	21																		
$80 < x \leq 100$	10	31																		
$100 < x \leq 120$	4	35																		
2.3	<p style="text-align: center;"><b>OGIVE</b></p>  <p style="text-align: center;"><b>Commission earned (in thousands of Rands)</b></p>	<p>✓ grounded/geanker</p> <p>✓ upper limits/ boonste limiet</p> <p>✓ cum frequency / Kum frekwensie</p> <p>✓ shape/vorm</p> <p style="text-align: right;">(4)</p>																		
2.4	<p>No. of salesmen awarded bonuses: <math>35 - 26</math> = 9 salesmen</p>	<p>✓ accept (25 – 27)</p> <p>✓ accept (8 – 10)</p> <p style="text-align: right;">(2)</p>																		
2.5	<p>Estimated mean = <math>\frac{(30 \times 7) + (50 \times 6) + (70 \times 8) + (90 \times 10) + (110 \times 4)}{35}</math></p> <p style="margin-left: 40px;"><math>= \frac{2410}{35}</math></p> <p style="margin-left: 40px;"><math>= 68,86</math> thousand rand or R68 857,14</p> <p style="margin-left: 40px;"><math>=</math> R69 000 or 69 thousand rand</p>	<p>✓ top line using midpts &amp; freq</p> <p>✓ 2410</p> <p>✓ answer (nearest)</p> <p style="text-align: right;">(3)</p> <p style="text-align: right;"><b>[12]</b></p>																		


**QUESTION/VRAAG 3**



3.1	$m_{CD} = \frac{-3 - (-5)}{4 - 0}$ $= \frac{-3 + 5}{4 - 0}$ $= \frac{1}{2}$	✓ substitution of C & D  ✓ answer  (2)
3.2	$m_{AD} = \frac{-5 - 3}{0 - (-4)}$ $= -2$ $m_{CD} \times m_{AD} = \frac{1}{2} \times -2$ $= -1$ $\therefore AD \perp DC$	✓ substitution of A & D ✓ $m_{AD} = -2$  ✓ product = -1  (3)
3.3	$AB = \sqrt{(3 + 4)^2 + (4 - 3)^2} = \sqrt{50} = 5\sqrt{2}$ $BC = \sqrt{(4 - 3)^2 + (-3 - 4)^2} = 5\sqrt{2}$ $AB = BC$ $\therefore \Delta ABC \text{ is an isosceles triangle/'n gelykbenige driehoek}$	✓ correct substitution ✓ length of AB  ✓ correct substitution ✓ length of BC  (4)

<p>3.4</p>	$m_{CD} = m_{BF} = \frac{1}{2}$ <p style="text-align: center;">[BF    DC]</p> $4 = \frac{1}{2}(3) + c$ $c = \frac{5}{2}$ $y = \frac{1}{2}x + \frac{5}{2}$ <p style="text-align: center;"><b>OR/OF</b></p> $y - 4 = \frac{1}{2}(x - 3)$ $y - 4 = \frac{1}{2}x - 1\frac{1}{2}$ $y = \frac{1}{2}x + 2\frac{1}{2}$	<p>✓ <math>m_{BF} = \frac{1}{2}</math></p> <p>✓ substitution of B(3 ; 4)</p> <p>✓ equation (3)</p>
<p>3.5</p>	$\tan \alpha = -2$ $\therefore \alpha = 116,57^\circ$ $\alpha = 90^\circ + \theta$ <p style="text-align: center;">[ext <math>\angle \Delta</math>]</p> $\therefore \theta = 26,57^\circ$ <p><b>OR/OF</b></p> $\tan \alpha = -2 \text{ OR } m_{AD} = -2$ $\therefore \tan \theta = \frac{1}{2}$ $\therefore \theta = 26,57^\circ$ <p><b>OR/OF</b></p> <p>Inclination of DE is <math>\beta</math>:</p> $\tan \beta = \frac{1}{2}$ $\therefore \beta = 26,57^\circ$ $\therefore \hat{ODE} = 63,43^\circ$ $\therefore \theta = 90^\circ - 63,43^\circ$ $= 26,57^\circ$	<p>✓ <math>\tan \alpha = -2</math></p> <p>✓ <math>\alpha = 116,57^\circ</math></p> <p>✓ <math>\theta = 26,57^\circ</math> (3)</p> <p>✓ <math>\tan \alpha = -2</math></p> <p>✓ <math>\tan \theta = \frac{1}{2}</math></p> <p>✓ <math>\theta = 26,57^\circ</math> (3)</p> <p>✓ <math>\beta = 26,57^\circ</math></p> <p>✓ <math>\hat{ODE} = 63,43^\circ</math></p> <p>✓ <math>\theta = 26,57^\circ</math> (3)</p>
<p>3.6</p>	$x^2 + y^2 = r^2$ $(4)^2 + (-3)^2 = 25$ $x^2 + y^2 = 25$	<p>✓ <math>r^2 = 25</math></p> <p>✓ equation (2)</p> <p style="text-align: right;"><b>[17]</b></p>

**QUESTION/VRAAG 4**



4.1	$N\left(\frac{1+(-3)}{2}; \frac{4+(-2)}{2}\right)$ <p>N(-1; 1) is the centre of the circle</p>	✓ substitution M & P ✓ x-value of N ✓ y-value of N (3)
4.2	$r = \sqrt{(1-(-1))^2 + (4-1)^2}$ $r = \sqrt{13} = \text{radius}$ $(x+1)^2 + (y-1)^2 = 13$ <p><b>OR/OR</b></p> $r = \sqrt{(-3-(-1))^2 + (-2-1)^2}$ $r = \sqrt{13} = \text{radius}$ $(x+1)^2 + (y-1)^2 = 13$	✓ substitution N & P ✓ $r = \sqrt{13}$ ✓ LHS of eq ✓ RHS of eq (4)
		✓ substitution N & M ✓ $r = \sqrt{13}$ ✓ LHS of eq ✓ RHS of eq (4)


<p>4.3</p>	<p><math>m_{NM} \times m_{MR} = -1</math> [radius <math>\perp</math> tangent/raakklyn]  <math>m_{NM} = \frac{1 - (-2)}{-1 - (-3)}</math> OR <math>m_{PM} = \frac{4 - (-2)}{1 - (-3)}</math>  <math>= \frac{3}{2}</math> <math>= \frac{3}{2}</math>  <math>m_{MR} = -\frac{2}{3}</math>  <math>y - y_1 = -\frac{2}{3}(x - x_1)</math> OR/OF <math>y = -\frac{2}{3}x + c</math>  <math>y + 2 = -\frac{2}{3}(x + 3)</math> OR/OF <math>-2 = -\frac{2}{3}(-3) + c</math>  <math>y = -\frac{2}{3}x - 4</math></p>	<p>✓ correct substitution                  ✓ <math>m_{NM}</math>                  ✓ <math>m_{MR}</math>                  ✓ substitution of <math>m_{MR}</math> &amp; <math>(-3; -2)</math>                  ✓ equation                  (5)</p>
<p>4.4</p>	<p>Symmetry of a kite: <math>S(-3; 4)</math>                  OR/OF  <math>\hat{P}SM = 90^\circ</math> [<math>\angle</math> in semi circle]  <math>PS \perp SM</math>  <math>\therefore S(-3; 4)</math>                  OR/OF  <math>(NS)^2 = (\text{radius})^2</math>  <math>(-3+1)^2 + (y-1)^2 = 13</math>  <math>(y-1)^2 = 9</math>  <math>y-1 = \pm 3</math>  <math>y = 4</math> OR <math>y \neq -2</math>  <math>\therefore S(-3; 4)</math></p>	<p>✓ x-value of S                  ✓ y-value of S                  (2)                  ✓ x-value of S                  ✓ y-value of S                  (2)                  ✓ x-value of S                  ✓ y-value of S                  (2)</p>
<p>4.5</p>	<p><math>(SR)^2 = (RM)^2</math> ...Tangents from common pt/rklyne v dies punt  <math>(x+3)^2 + (y-4)^2 = (x+3)^2 + (y+2)^2</math>  <math>y^2 - 8y + 16 = y^2 + 4y + 4</math>  <math>-12y = -12</math>  <math>y = 1</math>  <math>\frac{2}{3}x = -4 - 1</math> or <math>1 = -\frac{2}{3}x - 4</math>  <math>x = -\frac{15}{2}</math> <math>x = -7\frac{1}{2}</math>  <math>\therefore R\left(-7\frac{1}{2}; 1\right)</math>                  OR/OF</p>	<p>✓ equating lengths                  ✓ simplification                  ✓ y-value of R                  ✓ x-value of R                  (4)</p>



	$R(x;1)$ $\therefore 1 = -\frac{2}{3}x - 4$ $5 = -\frac{2}{3}x$ $x = -\frac{15}{2}$ $\therefore R\left(-\frac{15}{2};1\right)$  <b>OR/OF</b> $m_{NS} = \frac{1-4}{-1+3} = -\frac{3}{2}$ $\therefore m_{RS} = \frac{2}{3}$ $y-4 = \frac{2}{3}(x+3)$ $y = \frac{2}{3}x + 6$ $-\frac{2}{3}x - 4 = \frac{2}{3}x + 6$ $x = -7\frac{1}{2}$ $y = \frac{2}{3}\left(-\frac{15}{2}\right) + 6 = 1$ $\therefore R\left(-\frac{15}{2};1\right)$	$\checkmark y_R = 1$ $\checkmark$ horizontal line OR $R$ lies on $y = 1$ $\checkmark$ equating  $\checkmark$ $x$ -value of $R$ $(x < -4,6)$      $\checkmark y = \frac{2}{3}x + 6$ $\checkmark$ equating $\checkmark$ $x$ -value of $R$ $(x < -4,6)$ $\checkmark$ $y$ -value of $R$	(4)
4.6	$RS = \sqrt{(-3+7,5)^2 + (4-1)^2}$ OR/OF $RM = \sqrt{(-3+7,5)^2 + (-2-1)^2}$ $RS = \frac{3\sqrt{13}}{2} = 5,41$ area of $RSNM = 2$ area of $\Delta RSN$ $= 2\left(\frac{1}{2}\right)(\sqrt{13})\left(\frac{3\sqrt{13}}{2}\right)$ $= \frac{39}{2}$ <b>OR/OF</b> 19,5 square units  <b>OR/OF</b>	$\checkmark$ <b>RS OR RM</b> $\checkmark$ method $\checkmark \sqrt{13}$ and $\left(\frac{3\sqrt{13}}{2}\right)$ $\checkmark$ answer   $\checkmark$ method $\checkmark$ $MS = 6$ $\checkmark$ $RN = 6,5$ $\checkmark$ answer	(4)

	<p>area RSNM = <math>\frac{1}{2}(MS \times RN)</math> (area of a kite/<i>opp v vlieër</i>)</p> $= \frac{1}{2}(6)(6,5)$ $= \frac{39}{2} \text{ OR } 19,5 \text{ square units}$ <p><b>OR/OF</b></p> $RS = \sqrt{(-3 + 7,5)^2 + (4 - 1)^2} \text{ OR/OF } RM = \sqrt{(-3 + 7,5)^2 + (-2 - 1)^2}$ $RS = \frac{3\sqrt{13}}{2} \text{ or } 5,41$ $\text{area of } \Delta RSN = \left(\frac{1}{2}\right)(\sqrt{13})\left(\frac{3\sqrt{13}}{2}\right)$ $= \frac{39}{4} \text{ OR/OF } 9,75 \text{ square units}$ <p>area of RSNM = 2area of <math>\Delta RSN</math></p> $= \frac{39}{2} \text{ OR/OF } 19,5 \text{ square units}$ <p><b>OR/OF</b></p> <p>SM = 6</p> <p>area of RSNM = Area of <math>\Delta SMN</math> + Area of <math>\Delta RSM</math></p> $= \frac{1}{2}(6)(1) + \frac{1}{2}(6)\left(5\frac{1}{2}\right)$ $= 3 + 16\frac{1}{2}$ $= 19\frac{1}{2}$	<p>(4)</p> <p>✓ RS <b>OR</b> RM</p> $\checkmark \left(\frac{1}{2}\right)\sqrt{13}\left(\frac{3\sqrt{13}}{2}\right)$ <p>✓ method ✓ answer</p> <p>(4)</p> <p>✓ method</p> <p>✓ MS = 6</p> <p>✓ <math>h = 1</math> &amp; <math>5\frac{1}{2}</math></p> <p>✓ answer</p> <p>(4)</p>
		<p>[22]</p>


**QUESTION/VRAAG 5**

<p>5.1.1</p>	$\tan A = \frac{\sin A}{\cos A}$ $= \frac{2p}{p}$ $= 2$ <p><b>OR/OF</b></p> $\tan A = \frac{2p}{p}$ $= 2$ <div style="text-align: center;">  <p><math>(p ; 2p)</math></p> </div>	<p>✓ identity</p> <p>✓ value of tan A (2)</p> <p>✓ <math>\frac{y}{x}</math></p> <p>✓ value of tan A (2)</p>
<p>5.1.2</p>	$\sin^2 A + \cos^2 A = 1$ $(2p)^2 + p^2 = 1$ $4p^2 + p^2 = 1$ $5p^2 = 1$ $p^2 = \frac{1}{5}$ $\therefore p = -\frac{1}{\sqrt{5}}$	<p>✓ <math>(2p)^2 + p^2 = 1</math></p> <p>✓ simplification of LHS</p> <p>✓ answer (3)</p>
<p>5.2</p>	$2 \sin^2 x - 5 \sin x + 2 = 0$ $(2 \sin x - 1)(\sin x - 2) = 0$ $\sin x = \frac{1}{2} \text{ or } \sin x = 2(\text{no solution})$ <p>ref <math>\angle = 30^\circ</math></p> $\therefore x = 30^\circ + k.360^\circ \text{ or } x = 150^\circ + k.360^\circ ; k \in Z$	<p>✓ factors or formula</p> <p>✓ both equations</p> <p>✓ no solution/geen opl</p> <p>✓ <math>30^\circ + k.360^\circ</math></p> <p>✓ <math>150^\circ + k.360^\circ ;</math></p> <p>✓ <math>k \in Z</math> (6)</p>
<p>5.3.1</p>	$\sin(x + 300^\circ) = \sin x \cos 300^\circ + \cos x \sin 300^\circ$	<p>✓ expansion/uitbreiding (1)</p>
<p>5.3.2</p>	$\sin(x + 300^\circ) - \cos(x - 150^\circ)$ $= \sin x \cos 300^\circ + \cos x \sin 300^\circ - (\cos x \cos 150^\circ + \sin x \sin 150^\circ)$ $= \sin x \cos 60^\circ - \cos x \sin 60^\circ - (-\cos x \cos 30^\circ + \sin x \sin 30^\circ)$ $= \sin x \cos 60^\circ - \cos x \sin 60^\circ + \cos x \cos 30^\circ - \sin x \sin 30^\circ$ $= \frac{1}{2} \sin x - \frac{\sqrt{3}}{2} \cos x + \frac{\sqrt{3}}{2} \cos x - \frac{1}{2} \sin x$ $= 0$ <p><b>OR/OF</b></p>	<p>✓ 2<sup>nd</sup> expansion/ 2de uitbreiding</p> <p>✓✓ reduction/reduksie</p> <p>✓ special angle values/ spesiale hoekwaardes</p> <p>✓ answer (5)</p>

	$\begin{aligned} & \sin(x + 300^\circ) - \cos(x - 150^\circ) \\ &= \sin x \cos 300^\circ + \cos x \sin 300^\circ - (\cos x \cos 150^\circ + \sin x \sin 150^\circ) \\ &= \sin x \cos 60^\circ - \cos x \sin 60^\circ - (-\cos x \cos 30^\circ + \sin x \sin 30^\circ) \\ &= \sin x \cos 60^\circ - \cos x \sin 60^\circ + \cos x \cos 30^\circ - \sin x \sin 30^\circ \\ &= \sin x \sin 30^\circ - \cos x \sin 60^\circ + \cos x \sin 60^\circ - \sin x \sin 30^\circ \\ &= 0 \end{aligned}$	<p>✓ 2<sup>nd</sup> expansion/ 2de uitbreiding ✓✓ reduction/reduksie  ✓ co-ratios / ko-verh ✓ answer (5)</p>
<p>5.4</p>	<p>Consider: <math>\frac{\tan x + 1}{\sin x \tan x + \cos x} = \sin x + \cos x</math></p> $\text{LHS} = \frac{\left(\frac{\sin x}{\cos x} + 1\right)}{\left(\sin x \cdot \frac{\sin x}{\cos x} + \cos x\right)} = \frac{\left(\frac{\sin x + \cos x}{\cos x}\right)}{\left(\frac{\sin^2 x + \cos^2 x}{\cos x}\right)}$ $= \frac{\sin x + \cos x}{\frac{\cos x}{1}}$ $= \frac{\sin x + \cos x}{\cos x} \times \frac{\cos x}{1}$ $= \sin x + \cos x$ <p>= RHS</p> <p><b>OR/OF</b></p> $\text{LHS} = \frac{\left(\frac{\sin x}{\cos x} + 1\right)}{\left(\sin x \cdot \frac{\sin x}{\cos x} + \cos x\right)} = \frac{\left(\frac{\sin x + \cos x}{\cos x}\right)}{\left(\frac{\sin^2 x + \cos^2 x}{\cos x}\right)}$ $= \frac{\left(\frac{\sin x}{\cos x} + 1\right)}{\frac{1}{\cos x}}$ $= \left(\frac{\sin x}{\cos x} + 1\right) \times \frac{\cos x}{1}$ $= \sin x + \cos x$ <p>= RHS</p>	<p>✓ identity of tan x ✓ <math>\frac{\sin x + \cos x}{\cos x}</math> ✓ <math>\frac{\sin^2 x + \cos^2 x}{\cos x}</math>  ✓ <math>\sin^2 x + \cos^2 x = 1</math>  ✓ simplify (5)</p> <p>✓ identity of tan x ✓ <math>\frac{\sin^2 x + \cos^2 x}{\cos x}</math>  ✓ <math>\sin^2 x + \cos^2 x = 1</math>  ✓ simplify ✓ multiplication (5)</p>
<p>5.5.1</p>	$\begin{aligned} & (\sqrt{1+k})^2 = (\sin x + \cos x)^2 \\ & 1+k = \sin^2 x + 2 \sin x \cos x + \cos^2 x \\ & 1+k = 1 + \sin 2x \\ & k = \sin 2x \end{aligned}$	<p>✓ square both sides ✓ <math>\sin^2 x + \cos^2 x = 1</math> ✓ sin 2x (3)</p>


5.5.2	<p>From 5.5.1</p> $\sin x + \cos x = \sqrt{1 + \sin 2x}$ $\therefore \text{max value: } \sin x + \cos x = \sqrt{1+1}$ $= \sqrt{2}$ <p><b>OR/OF</b></p> <p>Maximum value of <math>1 + \sin 2x = 1 + 1</math></p> $= 2$ $\therefore \text{maximum value of } \sin x + \cos x = \sqrt{2}$ <p><b>OR/OF</b></p> $(\sin x + \cos x)^2 = \sin^2 x + 2 \sin x \cos x + \cos^2 x$ $= 1 + \sin 2x$ $\therefore \text{max value } (\sin x + \cos x)^2 = 1 + 1 = 2$ $\therefore \text{max value } \sin x + \cos x = \sqrt{2}$	<p>✓ max of <math>\sin 2x = 1</math></p> <p>✓ answer (2)</p> <p>✓ max of <math>\sin 2x = 1</math></p> <p>✓ answer (2)</p> <p>✓ max of <math>\sin 2x = 1</math></p> <p>✓ answer (2)</p>
		<b>[27]</b>

**QUESTION/VRAAG 6**



6.1	Period = $180^\circ$	✓ answer (1)
6.2	$-75^\circ$	✓ answer (1)
6.3	$\sin 2x \leq \frac{1}{\sqrt{2}} \cos x + \frac{1}{\sqrt{2}} \sin x$ $\sin 2x \leq \cos 45^\circ \cdot \cos x + \sin 45^\circ \cdot \sin x$ $\sin 2x \leq \cos(x - 45^\circ)$ $x \in [-75^\circ ; 165^\circ]$	✓ $\cos 45^\circ \cdot \cos x + \sin 45^\circ \cdot \sin x$ ✓ $\cos(x - 45^\circ)$ ✓ ✓ answer (4)
		<b>[6]</b>

**QUESTION/VRAAG 7**



Figure/Figuur (i)






Figure / Figuur (ii)

7.1	$KC = 6 \text{ cm}$	✓ answer (1)
7.2	Let P be the point of intersection of KL and CB  $\frac{KP}{KC} = \sin 60^\circ$ $KP = 6 \sin 60^\circ$ $KP = 3\sqrt{3}$ or 5,20 $\therefore KL = 8 + 3\sqrt{3}$ or 13,20 cm	 ✓ trig ratio ✓ length of KP ✓ answer (3)
7.3	$DK^2 = 6^2 + 12^2$ $DK = \sqrt{180}$ or $6\sqrt{5}$ or 13,42 cm $\frac{\sin \hat{KDL}}{KL} = \frac{\sin \hat{DLK}}{DK}$ $\frac{\sin \hat{KDL}}{\sin \hat{DLK}} = \frac{KL}{DK}$ $= \frac{8 + 3\sqrt{3}}{6\sqrt{5}}$ or $\frac{13,20}{13,42}$ or 0,98	✓ $DK = 6\sqrt{5}$ ✓ use of sine rule ✓ $\frac{\sin \hat{KDL}}{\sin \hat{DLK}} = \frac{KL}{DK}$ ✓ answer (4)

**[8]**


**QUESTION/VRAAG 8**



8.1	$\hat{L} = 100^\circ$ [ ext $\angle$ cyclic quad = int opp $\angle$ / <i>buite <math>\angle</math> kdvh = tos <math>\angle</math></i> <b>OR/OF</b> $\hat{N}_1 = 80^\circ$ [ $\angle$ s on straight line] $\hat{L} = 100^\circ$ [opp $\angle$ s of cyclic quad]	$\checkmark$ S $\checkmark$ R (2)  $\checkmark$ S $\checkmark$ R (2)
8.2	$\hat{N}_1 = 80^\circ$ [ $\angle$ s on straight line/ $\angle$ e op reguitlyn ] $\therefore \hat{O}_1 = 160^\circ$ [ $\angle$ at centre = $2 \times \angle$ at circumference/midpts $\angle = 2$ omtreks $\angle$ ] <b>OR/OF</b> reflex $\hat{K}\hat{O}\hat{M} = 200^\circ$ [ $\angle$ at centre = $2 \times \angle$ at circumference/midpts $\angle = 2 \times$ omtreks $\angle$ ] $\therefore \hat{O}_1 = 160^\circ$ [ $\angle$ s around a pt/ $\angle$ e om 'n pt]	$\checkmark$ S $\checkmark$ S $\checkmark$ R (3)  $\checkmark$ S $\checkmark$ R $\checkmark$ S (3)
8.3	$\hat{M}_1 = 360^\circ - (100^\circ + 55^\circ + 160^\circ)$ [sum $\angle$ s of quad/som $\angle$ e v vierhoek] $\therefore \hat{M}_1 = 45^\circ$	$\checkmark$ S $\checkmark$ S (2) <b>[7]</b>




**QUESTION/VRAAG 9**



9.1.1	$\angle$ in semi-circle/ $\angle$ in halfsirkel	✓ answer (1)
9.1.2	<b>Opp</b> $\angle$ s of quad = $180^\circ$ / <i>Teenoorst</i> $\angle$ e v vierhoek = $180^\circ$	✓ answer (1)
9.2.1	OF $\perp$ AC [line from centre bisects chord/ <i>lyn v midpt halv kd</i> ] $\therefore$ AC $\parallel$ GO [co-interior/ <i>ko-binne</i> $\angle$ s = $180^\circ$ <b>OR/OF</b> corresp/ <i>ooreenkomstige</i> $\angle$ s =]	✓ S ✓ R ✓ R (3)
9.2.2	$\hat{G}_1 = \hat{A}_2$ [corresp/ <i>ooreenk</i> $\angle$ s; AC $\parallel$ GO] $\hat{A}_2 = \hat{B}_1$ [ $\angle$ s in same segment/ <i>\angle</i> e in dies segment] $\therefore \hat{G}_1 = \hat{B}_1$ <b>OR/OF</b> $\hat{G}_1 = \hat{B}_2$ [ext $\angle$ cyclic quad/ <i>buite</i> $\angle$ koordevh] but $\triangle ABF \cong \triangle CBF$ [s, $\angle$ , s] $\therefore \hat{B}_2 = \hat{B}_1$ $\therefore \hat{G}_1 = \hat{B}_1$	✓ S ✓ R ✓ S ✓ R (4) ✓ S ✓ R ✓ R ✓ S (4)
9.3	OF : FB = 3 : 2 $\therefore$ DO = 5k and DF = 8k $\therefore \frac{DG}{DA} = \frac{DO}{DF} = \frac{r}{\frac{8}{5}r}$ $\therefore \frac{DG}{DA} = \frac{5}{8}$ <b>OR/OF</b> DB = 2r DF = $2r - \frac{2}{5}r = \frac{8}{5}r$ [line $\parallel$ side of $\triangle$ / <i>lyn <math>\parallel</math> sy v <math>\triangle</math></i> ]	✓ S ✓ R ✓ S (3)
		<b>[12]</b>

**QUESTION/VRAAG 10**




10.1	Tangent-chord theorem	✓ R	(1)
10.2.1	$\hat{A}_2 + \hat{A}_3 = \hat{B}_1 + \hat{B}_2$ [∠ <sup>s</sup> opp = sides/∠eteenoor = sye] $\hat{S}_3 = \hat{B}_1 + \hat{B}_2$ [ext ∠ cyclic quad/buite ∠ koordevh] $\therefore \hat{S}_3 = \hat{A}_2 + \hat{A}_3$ $\therefore AB \parallel ST$ [corresp/ooreenk ∠ <sup>s</sup> =]	✓ S ✓ R ✓ S ✓ R ✓ R	(5)
	<b>OR/OF</b>		
	$\hat{R}\hat{T}\hat{S} = \hat{B}\hat{A}\hat{S}$ [ext ∠ cyclic quad/buite ∠ koordevh] $\hat{B}\hat{A}\hat{S} = \hat{A}\hat{B}\hat{T}$ [∠ <sup>s</sup> opp = sides/∠eteenoor = sye] $\therefore \hat{R}\hat{T}\hat{S} = \hat{A}\hat{B}\hat{T}$ $\therefore AB \parallel ST$ [corresp/ooreenk ∠ <sup>s</sup> =]	✓ S ✓ R ✓ S ✓ R ✓ R	(5)

10.2.2	$\hat{B}_2 = x$ [tan chord theorem/raakl – koordst] $x + \hat{T}_4 = \hat{B}_1 + \hat{B}_2$ [corresp/ooreenk $\angle^s$ ; AB // ST] $\therefore \hat{T}_4 = \hat{B}_1$ $\hat{B}_1 = \hat{A}_1$ [tan chord theorem/raakl – koordst] $\therefore \hat{T}_4 = \hat{A}_1$	✓ S ✓ R ✓ S ✓ R ✓ R (5)
10.2.3	$\hat{T}_4 = \hat{A}_1$ [proven/bewys in 10.2.2] $\therefore$ RTAP is a cyclic quadrilateral [line subtends = $\angle^s$ ] <i>Is 'n koordevierhoek [lyn onderspan = <math>\angle e</math>]</i>	✓ S ✓ R (2)
		<b>[13]</b>


**QUESTION/VRAAG 11**

11.1



<p>11.1</p>	<p>Constr: On sides AB and AC of <math>\triangle ABC</math>, mark points G and H respectively such that <math>AG = DE</math> and <math>AH = DF</math>. Draw GH/Merk punt G en H op sy AB en AC van <math>\triangle ABC</math> onderskeidelik af sodanig dat <math>AG = DE</math> en <math>AH = DF</math>. Trek GH.</p> <p>Proof/Bewys:</p> <p><math>\triangle AGH \equiv \triangle DEF</math> [s, <math>\angle</math>, s]</p> <p><math>\therefore \hat{A}GH = \hat{E}</math></p> <p><math>= \hat{B}</math> [<math>\hat{B} = \hat{E}</math>, given/gegee]</p> <p><math>\therefore GH \parallel BC</math> [corresp/ooreenk <math>\angle^s =</math>]</p> <p><math>\therefore \frac{AG}{AB} = \frac{AH}{AC}</math> [line <math>\parallel</math> side of <math>\triangle</math> / lyn <math>\parallel</math> sye v <math>\triangle</math>]</p> <p><math>\therefore \frac{DE}{AB} = \frac{DF}{AC}</math> [constr/konstruksie]</p>	<p>✓ construction/ konstruksie</p> <p>✓ S/R</p> <p>✓ S</p> <p>✓ S /R</p> <p>✓ S ✓ R</p> <p>(6)</p>
-------------	---	--

11.2



11.2.1(a)	$AP = PC$ [diag $\parallel^m$ bisect each other/ <i>hoekl</i> $\parallel^m$ <i>halveer mekaar</i> ] But $TP = PS$ [given/ <i>gegee</i> ] $AP - TP = PC - PS$ $\therefore AT = SC$	$\checkmark$ S $\checkmark$ S OR S 2)
11.2.1(b)	In $\Delta PSR$ and $\Delta PBA$ : $\hat{P}_1 = \hat{P}_3$ [vertically opp $\angle^s$ / <i>regoorst</i> $\angle e$ ] $\hat{B}_1 = \hat{S}_1$ [ $\angle^s$ in same segment / $\angle e$ in <i>dies segment</i> ] $\therefore \Delta PSR \parallel \Delta PBA$ [ $\angle, \angle, \angle$ ]  <b>OR/OF</b> In $\Delta PSR$ and $\Delta PBA$ : $\hat{P}_1 = \hat{P}_3$ [vertically opp $\angle^s$ / <i>regoorst</i> $\angle e$ ] $\hat{B}_1 = \hat{S}_1$ [ $\angle^s$ in same segment / $\angle e$ in <i>dies segment</i> ] $\hat{A}_2 = \hat{R}_1$ [sum $\angle^s \Delta$ / <i>som</i> $\angle e \Delta$ ] $\therefore \Delta PSR \parallel \Delta PBA$ [ $\angle, \angle, \angle$ ]	$\checkmark$ S $\checkmark$ R $\checkmark$ S $\checkmark$ R $\checkmark$ R (5)  $\checkmark$ S $\checkmark$ R $\checkmark$ S $\checkmark$ R $\checkmark$ S (5)

<p>11.2.2(a)</p>	$\frac{PR}{PA} = \frac{PS}{PB} \quad [    \Delta s]$ $\therefore \frac{PR}{PA} = \frac{TR}{AD} = \frac{PS}{PB} \quad \left[ \text{given } \frac{PR}{PA} = \frac{TR}{AD} \right]$ $\therefore \frac{PR}{PA} = \frac{TR}{AD} = \frac{TP}{PD} \quad [PS = TP; PB = PD]$ $\therefore \Delta RPT \parallel \Delta APD \quad [\text{sides of } \Delta \text{ in prop/sye v } \Delta \text{ in dies verhouding}]$	<p>✓ S (all 3 ratios)                  ✓ S                  ✓ R                  (3)</p>
<p>11.2.2(b)</p>	$\hat{T}_1 = \hat{D}_2 \quad [    \Delta s]$ <p>∴ ATRD is a cyclic quad [converse: ext ∠ of cyclic quad/  <i>Omgekeerde buite ∠ v koordevh</i>]</p>	<p>✓ S                  ✓ R                  (2)</p>
		<p><b>[18]</b></p>

**TOTAL/TOTAAL: 150**